

Ankle Rehabilitation

Non-surgical

Phase I

- Modalities:
 - prn for pain and swelling (ice, IFC)
 - Consider contrast bath if significant edema
- Brace: as directed by physician
- ROM:
 - PROM and stretches
 - AROM in pain-free range
 - Join mobs to talocrural, subtalar, inter-metatarsal and MTP joints
- Exercises/Strengthening:
 - Isometrics
 - Knee/hip strengthening – within any WB precautions

Phase II

- Modalities: Continue prn
- Gait:
 - WB as directed by physician
 - Continue weight-shifting, progressing as tolerated
- ROM: Continue as in phase I
- Strengthening/Balance:
 - Light manual resistive exercises
 - Progress to theraband as tolerated
 - Knee/hip strengthening, progressing to closed-chain as tolerated – within any WB precautions

Phase III

- Modalities: Continue prn
- Gait: WBAT
- Brace: D/C'd
- ROM: Continue as II but more aggressive
- Strengthening/Balance:
 - BAPS
 - Balance board
 - Progress from double to single leg balance activities
 - Progress with closed-chain strengthening

Initiation Date: 01-01-2005 Revised Date: 06-11-14

Phone (815) 484-6990 * Fax (815) 484-6961



Phase IV

- Advance to running and agility drills, plyometrics, sport-specific activities as tolerated
- Testing: Less than 25% deficit for non-athletes, less than 20% for athletes

****Progress from one phase to the next once patient has achieved all goals in previous phase with no increased pain levels****

Adapted From:

- 1) Brotzman SB, Wilk KE. Clinical Orthopedic Rehabilitation. 2nd Ed. Philadelphia: Mosby; 2003
- 2) Zachazewski JE et al. Athletic Injuries and Rehabilitation. Philadelphia: WB Saunders CO; 1996