



---

## **GEOFFREY S. VAN THIEL, MD/MBA**

*Assistant Professor - Rush University Medical Center*

*Team Physician - US National Soccer Teams*

*Team Physician - Chicago Blackhawks Medical Network - Ice Hogs*

www.VanThielMD.com - VanThielMD@orthoillinois.com

Fax: (815) 381-7489 – Phone: (815) 381-7365



### **SPORTS MEDICINE SURGERY – HIP ARTHROSCOPY**

*Follow us* on Twitter and Facebook: @VanThielMD & Facebook.com/VanThielMD

– Cutting edge information on the arthroscopic treatment of the hip, knee and shoulder –

---

## **REHABILITATION PROTOCOL**

### **ACL Reconstruction**

(Patellar Tendon, Hamstring, or Allografts)

#### **Precautions –**

##### ***Revision ACL Reconstructions***

Per specific physician recommendation, follow protocol until 12 weeks, then extend weeks 12 to 16 through to 5- to 6-month timeline, when patients can then begin running and progress to functional sports activities.

##### ***Meniscus Repair***

If a meniscus repair was performed the patient is to remain foot flat (25%) weightbearing until 4 weeks.

#### **Phase I (1 – 10 days post-op)**

- Wound care: Observe for signs of infection. OK to remove dressing on post-operative day 5 and begin showering. Keep covered until day 5. Cover incision with gauze and ace wrap.
- Weight Bearing: Weight bearing as tolerated with the brace locked in extension. If a meniscus repair was performed, then foot flat (25%) weight bearing until week 4.
- Modalities: prn for pain and inflammation (ice, IFC)
- Brace: Immobilizer or brace if prescribed (hinged brace locked in full extension) To be worn at all times including when sleeping.
- ROM: Goal: Minimum 0 – 90 degrees, not more than 120 degrees
  - Passive positional stretches for extension and flexion
  - Ankle AROM

### Phase II (10 days – 4 weeks post-op)

- Wound care: Observe for signs of infection and begin scar management techniques when incision is closed. OK to remove dressing on post-operative day 5 and begin showering. Keep covered until day 5. Cover incision with gauze and ace wrap.
- Weight Bearing: Weight bearing as tolerated with the brace unlocked. If a meniscus repair was performed, then foot flat (25%) weight bearing until week 4 with the brace unlocked.
- Brace: Hinged brace set 0 – 120 and unlocked for ambulation. On at all times except in PT clinic. Discontinue brace use at night.
- ROM: Goal: Minimum 0 – 90 degrees, not more than 120 degrees until 3 weeks, then gradually to full AROM.
  - Passive positional stretches and AROM for extension and flexion
  - Half revolutions on stationary bike and progress to full revolutions
  - Increase / maintain patellar mobility with emphasis on superior glide
- Strengthening:
  - No resisted open chain strengthening
  - Quad sets (open and closed chain multi angle)
  - SLR (eliminate extensor lag)
  - Emphasize closed chain activities for strengthening (step ups, light leg press etc.)
  - Proprioceptive activities added as soon as quad control allows.
  - Balance board bilateral in multiple planes
  - Single-leg balance eyes open/closed, variable surfaces
- Modalities:
  - NMES to quads if unable to perform quad sets and extensor lag with SLR
  - IFC and ice for pain and edema prn
  - sEMG neuromuscular re-education for quad sets
- Conditioning
  - Upper Body Cycle
  - Stationary bike with gradual progressive resistance

### Phase III (4 – 8 weeks post-op)

- Wound care: Continue scar mobs
- Brace: Gradually discontinue brace from week 4 to 6
- ROM:
  - Emphasize full extension
  - Full flexion by end of 8 weeks
  - Patellar mobility
  - Rectus femoris/ hip flexor stretches
- Strengthening:
  - Continue Phase II, adding resistance as tolerated
  - Stationary bike: increase resistance and some light intervals
  - Squats/leg press: bilateral to unilateral (0–60 degrees) with progressive resistance
  - Lunges (0–60 degrees)
  - Stairs: concentric and eccentric (not to exceed 60 degrees of knee flexion)
  - Calf raises: bilateral to unilateral
  - Rotational stability exercises: static lunge with lateral pulley repetitions
  - Sport cord resisted walking all four directions

- Treadmill walking all four directions
- Balance board: multiple planes, bilateral stance
- Ball toss to mini-tramp or wall in single-leg stance
- Core strengthening: supine and prone bridging, standing with pulleys
- Gait activities: cone obstacle courses at walking speeds in multiple planes
- Modalities:
  - Continue E-stim for re-ed or edema
  - sEMG to continue (for balance of VL to VMO or overall contraction)
  - Continue ice and IFC prn
- Conditioning:
  - Stepper (retro and / or forward)
  - Stationary bike
  - UBC
  - Pool if available
- Gait: Normalize gait pattern on level surfaces and progress to step-over-step pattern on stairs

#### **Phase IV (8 - 12 weeks post-op)**

- Wound care: Continue scar mobs
- ROM: Full ROM
- Strengthening:
  - Increase weights and reps of previous exercises
  - Squats/leg press: bilateral to unilateral (0–60 degrees) progressive resistance
  - Lunges (0–60 degrees)
  - Calf raises: bilateral to unilateral
  - Advance hamstring strengthening
  - Core strengthening
  - Combine strength and balance (e.g., ball toss to trampoline on balance board, mini-squat on balance board, Sport Cord cone weaves, contrakicks)
  - Advanced balance exercises (e.g., single-leg stance while reaching to cones on floor with hands or opposite foot, single-leg stance while pulling band laterally)
  - Lap swimming generally fine with exception of breaststroke; caution with deep squat push-off and no use of fins yet
  - Stationary bike intervals
- Modalities: continue prn

#### **Phase V (12 – 16 weeks)**

- Important – Focus on correct technique
  - Landing during exercises at low knee flexion angles (too close to extension)
  - Landing during exercises with genu varum/valgum (watch for dynamic valgus of knee and correct)
  - Landing and jumping with uninjured limb dominating effort
- Exercises
  - Elliptical trainer: forward and backward
  - Perturbation training\*: balance board, roller board, roller board with platform
  - Shuttle jumping: bilateral to alternating to unilateral, emphasis on landing form
  - Mini-tramp bouncing: bilateral to alternating to unilateral, emphasis on landing form

- Jogging in place with sport cord: pulling from variable directions
- Movement speed increases for all exercises
- Slide board exercises
- Aqua jogging

### Phase VI (16 – 24 weeks)

- Exercises
  - Progressive running program
    - Always begin with warmup on the stationary bike or elliptical for >10 minutes prior to initiation of running.
    - Patient should have no knee pain following run.
    - Week 1: Run: walk 30 seconds: 90 seconds every other day (qod) (10–15 minutes)
    - Week 2: Run: walk 60:60 qod (10–20 minutes)
    - Week 3: Run: walk 90:30 qod (15–20 minutes)
    - Week 4: Run: walk 90:30 3-4x/week (20–25 minutes)
    - Week 5: Run continuously 15–20 minutes 3–5x/week
  - Hop testing and training
    - Single-leg hop for distance: 80% minimum compared to nonsurgical side for running, 90% minimum for return to sport
    - Single-leg triple hop for distance: 80% for running, 90% for return to sport
    - Triple crossover hop for distance: 80% for running, 90% for return to sport
    - Timed 10-m single-leg hop: 80% for running, 90% for return to sport
    - Timed vertical hop test: 60 seconds with good form and steady rhythm considered passing
  - Vertical, horizontal jumping from double to single leg
  - Progressive plyometrics (e.g., box jumps, bounding, standing jumps, jumps in place, depth jumps, squat jumps, scissor jumps, jumping over barriers, skipping)
  - Speed and agility drills (e.g., T-test, line drills) (make these similar in movement to specific sport of athlete).
  - Cutting drills begin week 20
  - Progress to sport-specific drills week 20
  - Return to Sport at 6 months

Adapted From

1) Brotzman SB, Manske RC. Clinical Orthopedic Rehabilitation. 3rd Ed. Elsevier; 2011.