


## Heel Pain

Heel pain is most often caused by plantar fasciitis -- a condition that is sometimes also called heel spur syndrome when a spur is present. Heel pain may also be due to other causes, such as a stress fracture, tendonitis, arthritis, or nerve irritation. Because there are several potential causes, it is important to have heel pain properly diagnosed.

### What is Plantar Fasciitis?

Plantar fasciitis is an inflammation of the band of tissue (the plantar fascia) that extends from the heel to the toes. In this condition, the fascia first becomes irritated and then inflamed—resulting in heel pain.



### Symptoms of Plantar Fasciitis

The symptoms of plantar fasciitis are:

- Pain on the bottom of the heel
- Pain that is usually worse upon arising
- Pain that increases over a period of months

People with plantar fasciitis often describe the pain as worse when they get up in the morning or after they've been sitting for long periods of time. After a few minutes of walking the pain decreases, because walking stretches the fascia.

For some people the pain subsides but returns after spending long periods of time on their feet.

### Causes of Plantar Fasciitis

The most common cause of plantar fasciitis relates to faulty mechanics of the foot. For example, people who have problems with their arches—either overly flat feet or high-arched feet—are more prone to developing plantar fasciitis. People with tight Achilles tendons, also are more prone to develop plantar fasciitis.

Wearing non-supportive footwear on hard, flat surfaces puts abnormal strain on the plantar fascia and can also lead to plantar fasciitis. This is particularly evident when a person's job requires long hours on their feet. Obesity also contributes to plantar fasciitis.

## Diagnosis

To arrive at a diagnosis, Dr. Sorensen will obtain your medical history and examine your foot. Throughout this process Dr. Sorensen will rule out all the possible causes for your heel pain other than plantar fasciitis.

In addition, diagnostic imaging studies such as x-rays, or magnetic resonance imaging (MRI) may be used to distinguish the different types of heel pain. Sometimes heel spurs are found in patients with plantar fasciitis, but these are rarely a source of pain. When they are present, the condition may be diagnosed as plantar fasciitis/heel spur syndrome.

## Treatment Options

Treatment of plantar fasciitis begins with first-line strategies, which you can begin at home:

- **Stretching exercises.** Exercises that stretch out the calf muscles help ease pain and assist with recovery.
- **Avoid going barefoot.** When you walk without shoes, you put undue strain and stress on your plantar fascia.
- **Ice.** Putting an ice pack on your heel for 15 minutes several times a day helps reduce inflammation. Limit activities. Cut down on extended physical activities to give your heel a rest.
- **Shoe modifications.** Wearing supportive shoes that have good arch support and a slightly raised heel reduces stress on the plantar fascia. Your shoes should provide a comfortable environment for the foot.
- **Medications.** Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, may help reduce pain and inflammation.
- **Lose weight.** Extra pounds put extra stress on your plantar fascia.

If you still have pain after several weeks, Dr Sorensen may add one or more of these approaches:

- **Orthotic devices.** Custom orthotic devices that fit into your shoe help correct the underlying structural abnormalities causing the plantar fasciitis.
- **Injection therapy.** In some cases, corticosteroid injections are used to help reduce the inflammation and relieve pain.
- **Removable walking cast.** A removable walking cast may be used to keep your foot immobile for a few weeks to allow it to rest and heal.
- **Night splint.** Wearing a night splint allows you to maintain an extended stretch of the plantar fascia while sleeping. This may help reduce the morning pain experienced by some patients.
- **Physical therapy.** Exercises and other physical therapy measures may be used to help provide relief.

Although most patients with plantar fasciitis respond to non-surgical treatment, a small percentage of patients may require surgery. If, after several months of non-surgical treatment, you continue to have heel pain, surgery may be considered.

## Long-Term Care

No matter what kind of treatment you undergo for plantar fasciitis, the underlying causes that led to this condition may remain. Therefore, you will need to continue with preventive measures. Wearing supportive shoes, using custom orthotic devices and continuing stretching are the mainstay of long-term treatment for plantar fasciitis.

\*View Dr. Sorensen's Web MD article at: <http://emedicine.medscape.com/article/1236763-overview>

