

MEDICATIONS TO WITHHOLD BEFORE SURGERY

PLEASE STOP TAKING ANY OF THE MEDICATIONS LISTED BELOW 7 DAYS PRIOR TO YOUR SURGERY. This includes any aspirin or aspirin containing medication, blood thinning medication, and any non-steroidal anti-inflammatory (NSAIDs) medication. All of these medications thin your blood and could cause you to lose more blood than expected during surgery. You may resume your medication as instructed by your surgeon. Please refer to your postoperative instructions to see when you may resume your medication after surgery. If you have any questions or concerns regarding your medication, please contact your surgeon's office.

BRAND NAMES

Advil	Ecotrin
Aggrastat	Effient
Aggrenox	Eliquis
Aleve	Excedrin
Alka Seltzer	Feldene
Amigesic	Fiorinal
Anacin	Fish oil
Anaprox	Flector
Angiomax	Fragmin
Argatroban	Halfprin
Arixtra	Heparin
Arthrotec	Herbal Supplements
Asacol	Indocin
Ascriptin	Integrilin
Aspirin	Iprivask
Azulfidine	Kaopectate
Bayer	Lovenox
Brilinta	Mobic
Bufferin	Motrin
Cambia	Naprelan
Canasa	Naprosyn
Cataflam	Norgesic
Celebrex	Norgesic Forte
Cilostazol	Orudis
Clinoril	Pepto Bismol
Colazal	Persantine
Coumadin	Plavix
Daypro	Pletal
Dipentum	Pradaxa
Doan's	Relafen
Dolobid	ReoPro
	Salflex

Savaysa
Soma compound
Ticlid
Tolectin
Toradol
Treximet
Tricosal
Vicoprofen
Voltaren
Xarelto
Zipsor
Zontivity
Zorvolex

Enoxaparin
Eptifibatide
Etodolac
Fondaparinux
Ibuprofen
Indomethacin
Ketoprofen
Ketorolac tromethamine
Magnesium salicylate
Meclofenamate
Meloxicam
Mesalamine
Nabumetone
Naproxen
Olsalazine sodium
Oxaprozin
Piroxicam
Prasugrel
Rivaroxaban
Salsalate
Sulfaxalazine
Sulindac
Sumatriptan-naproxen
Ticagrelor
Ticlopidine HCL
Tirofiban
Tolmetin
Vorapaxar
Warfarin

GENERIC NAMES

Abciximab
Apixaban
Aspirin
Balsalazide
Bismuth subsalicylate
Bivalirudin
Celecoxib
Choline magnesium
Cilostazol
Clopidogrel
Dabigatran
Dalteparin
Desirudin
Diclofenac
Diflunisal
Dipyridamole
Edoxaban

FOR PATIENTS UNDERGOING FUSION:

1. Do not take any osteoporosis medications (Actonel, Atelvia, Binosto, Boniva, Evista, Fosamax, Prolia, Reclast, etc.) for one month before and three months after surgery, unless specifically discussed with your surgeon.
2. Do not take any oral rheumatoid arthritis medications (Arava, Azasan, Azulfidine, Imuran, Plaquenil, etc.) for two weeks before and two weeks after surgery, unless specifically discussed with your surgeon.
3. Do not take any injectable rheumatoid arthritis medications (Cimzia, Enbrel, Humira, Orencia, Remicade, Rheumatrex, Xeljanz, etc.) for one month before and one month after surgery, unless specifically discussed with your surgeon.